

Product datasheet for RC201467L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Glutathione S Transferase kappa 1 (GSTK1) (NM_015917) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Glutathione S Transferase kappa 1 (GSTK1) (NM_015917) Human Tagged ORF Clone Lentiviral

Particle

Symbol: Glutathione S Transferase kappa 1

Synonyms: GST; GST13; GST 13-13; GSTK1-1; hGSTK1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_015917

ORF Size: 678 bp

ORF Nucleotide

Sequence:

Domains:

The ORF insert of this clone is exactly the same as(RC201467).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 015917.1</u>

RefSeq Size: 1058 bp
RefSeq ORF: 681 bp
Locus ID: 373156
UniProt ID: Q9Y2Q3
Cytogenetics: 7q34

DSBA





Glutathione S Transferase kappa 1 (GSTK1) (NM_015917) Human Tagged ORF Clone Lentiviral Particle – RC201467L3V

Protein Pathways: Drug metabolism - cytochrome P450, Glutathione metabolism, Metabolism of xenobiotics by

cytochrome P450

MW: 25.5 kDa

Gene Summary: This gene encodes a member of the kappa class of the glutathione transferase superfamily of

enzymes that function in cellular detoxification. The encoded protein is localized to the peroxisome and catalyzes the conjugation of glutathione to a wide range of hydrophobic substates facilitating the removal of these compounds from cells. Alternative splicing results

in multiple transcript variants.[provided by RefSeq, Jan 2009]