

Product datasheet for RC201478L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

HP1 gamma (CBX3) (NM 007276) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: HP1 gamma (CBX3) (NM_007276) Human Tagged ORF Clone Lentiviral Particle

Symbol: HP1 gamma

Synonyms: HECH; HP1-GAMMA; HP1Hs-gamma

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_007276

ORF Size: 549 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC201478).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 007276.3

 RefSeq Size:
 2330 bp

 RefSeq ORF:
 552 bp

 Locus ID:
 11335

 UniProt ID:
 Q13185

 Cytogenetics:
 7p15.2

Domains: CHROMO

Protein Families: Druggable Genome, Transcription Factors





MW: 20.8 kDa

Gene Summary:

At the nuclear envelope, the nuclear lamina and heterochromatin are adjacent to the inner nuclear membrane. The protein encoded by this gene binds DNA and is a component of heterochromatin. This protein also can bind lamin B receptor, an integral membrane protein found in the inner nuclear membrane. The dual binding functions of the encoded protein may explain the association of heterochromatin with the inner nuclear membrane. This protein binds histone H3 tails methylated at Lys-9 sites. This protein is also recruited to sites of ultraviolet-induced DNA damage and double-strand breaks. Two transcript variants encoding the same protein but differing in the 5' UTR, have been found for this gene. [provided by RefSeq, Mar 2011]