


## Product datasheet for RC201499L4

### TRMT61B (NM\_017910) Human Tagged Lenti ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	TRMT61B (NM_017910) Human Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	TRMT61B
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC201499).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF.

ACCN: NM\_017910


ORF Size: 1431 bp



[View online »](#)

<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"> <li>1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
<b>RefSeq:</b>	<a href="#">NM_017910.2</a> , <a href="#">NP_060380.2</a>
<b>RefSeq Size:</b>	1856 bp
<b>RefSeq ORF:</b>	1434 bp
<b>Locus ID:</b>	55006
<b>UniProt ID:</b>	<a href="#">Q9BVS5</a>
<b>Cytogenetics:</b>	2p23.2
<b>Protein Families:</b>	Druggable Genome
<b>MW:</b>	53 kDa
<b>Gene Summary:</b>	<p>Methyltransferase that catalyzes the formation of N(1)-methyladenine at position 58 (m1A58) in various tRNAs in mitochondrion, including tRNA(Leu) (deciphering codons UUA or UUG), tRNA(Lys) and tRNA(Ser) (deciphering codons UCA, UCU, UCG or UCC) (PubMed:23097428). Catalyzes the formation of 1-methyladenosine at position 947 of mitochondrial 16S ribosomal RNA and this modification is most likely important for mitoribosomal structure and function (PubMed:27631568). In addition to tRNA N(1)-methyltransferase activity, also acts as a mRNA N(1)-methyltransferase by mediating methylation of adenosine residues at the N(1) position of MT-ND5 mRNA, leading to interfere with mitochondrial translation (PubMed:29107537). [UniProtKB/Swiss-Prot Function]</p>

## Product images:



Circular map for RC201499L4