

Product datasheet for RC201544L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Integrin Linked Kinase (ILK) (NM 004517) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Integrin Linked Kinase (ILK) (NM 004517) Human Tagged ORF Clone Lentiviral Particle

Symbol: Integrin Linked Kinase

Synonyms: HEL-S-28; ILK-1; ILK-2; P59; p59ILK

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM_004517

ORF Size: 1356 bp

ORF Nucleotide

The ODE

Sequence:

The ORF insert of this clone is exactly the same as(RC201544).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 004517.2

 RefSeq Size:
 1843 bp

 RefSeq ORF:
 1359 bp

 Locus ID:
 3611

 UniProt ID:
 Q13418

 Cytogenetics:
 11p15.4

Domains: pkinase, TyrKc, ANK, S_TKc

Protein Families: Druggable Genome, Protein Kinase





Integrin Linked Kinase (ILK) (NM_004517) Human Tagged ORF Clone Lentiviral Particle – RC201544L1V

Protein Pathways: Endometrial cancer, Focal adhesion, PPAR signaling pathway

MW: 51.4 kDa

Gene Summary: This gene encodes a protein with a kinase-like domain and four ankyrin-like repeats. The

encoded protein associates at the cell membrane with the cytoplasmic domain of beta integrins, where it regulates integrin-mediated signal transduction. Activity of this protein is important in the epithelial to mesenchymal transition, and over-expression of this gene is implicated in tumor growth and metastasis. Alternative splicing results in multiple transcript

variants. [provided by RefSeq, Jun 2013]