

Product datasheet for RC201555L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RAB13 (NM_002870) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: RAB13 (NM_002870) Human Tagged ORF Clone Lentiviral Particle

Symbol: RAB13 Synonyms: GIG4

Mammalian Cell Puromycin

Selection:

Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_002870

ORF Size: 609 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC201555).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 002870.2

 RefSeq Size:
 1235 bp

 RefSeq ORF:
 612 bp

 Locus ID:
 5872

 UniProt ID:
 P51153

 Cytogenetics:
 1q21.3

Domains: ras, RAN, RAS, RHO, RAB, ARF

Protein Families: Druggable Genome





RAB13 (NM_002870) Human Tagged ORF Clone Lentiviral Particle - RC201555L3V

Protein Pathways: Tight junction

MW: 22.8 kDa

Gene Summary: This gene is a member of the Rab family of small G proteins and plays a role in regulating

membrane trafficking between trans-Golgi network (TGN) and recycling endosomes (RE). The encoded protein is involved in the assembly of tight junctions, which are components of the apical junctional complex (AJC) of epithelial cells. The AJC plays a role in forming a barrier between luminal contents and the underlying tissue. Additional functions associated with the protein include endocytic recycling of occludin, regulation of epithelial cell scattering, neuronal regeneration and regulation of neurite outgrowth. Alternately spliced transcript variants have been observed for this gene. A pseudogene associated with this gene is located

on chromosome 12. [provided by RefSeq, Jan 2013]