

Product datasheet for RC201582

H2BC12 (NM_080593) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: H2BC12 (NM_080593) Human Tagged ORF Clone
Tag: Myc-DDK
Symbol: H2BC12
Synonyms: H2B/S; H2BFAii; H2BFT; H2BK; HIST1H2BK
Mammalian Cell Selection: Neomycin
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
ORF Nucleotide Sequence: >RC201582 ORF sequence
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGCCGGAACAGCGAAGTCCGCTCCCGCGCCCAAGAAGGGCTCGAAGAAAGCCGTGACTAAGGCGCAGA
AGAAGGACGGCAAGAAGCGCAAGCGCAGCCGCAAGGAGAGCTACTCCGTATACGTGTACAAGGTGCTGAA
GCAGGTCCACCCGACACCGGCATCTCTCTAAGGCCATGGGAATCATGAACTCCTTCGTCACGACATC
TTCGACGCATCGCGGTGAGGCTTCCCGCCTGGCGCATTACAACAAGCGCTCGACCATCACCTCCAGGG
AGATCCAGACGGCCGTGCGCCTGCTGCTGCCCGGGGAGTTGGCCAAGCACGCCGTGCCAGGGCACCAA
GGCCGTCACCAAGTACACCAGCGCTAAG

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC201582 protein sequence
Red=Cloning site Green=Tags(s)
MPEPAKSAPAPKKGSKKAVTKAQKKGKRRSRKESYSVYVYKVLKQVHPDTGISSKAMGIMNSFVNDI
FERIAGEASRLAHYNKRSTITSREIQTAVRLLLPGELAKHAVSEGTKAVTKYTSK

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6301_g04.zip

Restriction Sites: SgfI-MluI



[View online »](#)

Cloning Scheme:


ACCN: NM_080593

ORF Size: 378 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

- Reconstitution Method:**
1. Centrifuge at 5,000xg for 5min.
 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
 3. Close the tube and incubate for 10 minutes at room temperature.
 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_080593.2](#)

RefSeq Size: 863 bp

RefSeq ORF: 381 bp


Locus ID: 85236

UniProt ID: [O60814](#)


Cytogenetics: 6p22.1
Protein Pathways: Systemic lupus erythematosus
MW: 13.9 kDa

Gene Summary: Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a replication-dependent histone that is a member of the histone H2B family. The protein encoded is an antimicrobial protein with antibacterial and antifungal activity. Two transcripts that encode the same protein have been identified for this gene, which is found in the histone microcluster on chromosome 6p21.33. [provided by RefSeq, Aug 2015]

Product images:



Circular map for RC201582



Western blot validation of overexpression lysate (Cat# [LY409140]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC201582 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).