

Product datasheet for RC201603L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CDK9 (NM_001261) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CDK9 (NM_001261) Human Tagged ORF Clone Lentiviral Particle

Symbol: CDK9

Synonyms: C-2k; CDC2L4; CTK1; PITALRE; TAK

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_001261

 ORF Size:
 1116 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC201603).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 001261.2

 RefSeq Size:
 2472 bp

 RefSeq ORF:
 1119 bp

 Locus ID:
 1025

 UniProt ID:
 P50750

 Cytogenetics:
 9q34.11

Domains: pkinase, TyrKc, S_TKc

Protein Families: Druggable Genome, Protein Kinase, Transcription Factors





ORIGENE

MW: 42.8 kDa

Gene Summary:

The protein encoded by this gene is a member of the cyclin-dependent protein kinase (CDK) family. CDK family members are highly similar to the gene products of S. cerevisiae cdc28, and S. pombe cdc2, and known as important cell cycle regulators. This kinase was found to be a component of the multiprotein complex TAK/P-TEFb, which is an elongation factor for RNA polymerase II-directed transcription and functions by phosphorylating the C-terminal domain of the largest subunit of RNA polymerase II. This protein forms a complex with and is regulated by its regulatory subunit cyclin T or cyclin K. HIV-1 Tat protein was found to interact with this protein and cyclin T, which suggested a possible involvement of this protein in AIDS. [provided by RefSeq, Jul 2008]