

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC201606L2V

## NCOA4 (NM\_005437) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	NCOA4 (NM_005437) Human Tagged ORF Clone Lentiviral Particle
Symbol:	NCOA4
Synonyms:	ARA70; ELE1; PTC3; RFG
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
Tag:	mGFP
ACCN:	NM_005437
ORF Size:	1842 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC201606).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 005437.1</u>
RefSeq Size:	3502 bp
RefSeq ORF:	1845 bp
Locus ID:	8031
UniProt ID:	<u>Q13772</u>
Cytogenetics:	10q11.22
Protein Families:	Druggable Genome, Transcription Factors
Protein Pathways:	Pathways in cancer, Thyroid cancer



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	NCOA4 (NM_005437) Human Tagged ORF Clone Lentiviral Particle – RC201606L2V
MW:	69.7 kDa
Gene Summary:	This gene encodes an androgen receptor coactivator. The encoded protein interacts with the androgen receptor in a ligand-dependent manner to enhance its transcriptional activity. Chromosomal translocations between this gene and the ret tyrosine kinase gene, also located on chromosome 10, have been associated with papillary thyroid carcinoma. Alternatively spliced transcript variants have been described. Pseudogenes are present on chromosomes 4, 5, 10, and 14. [provided by RefSeq, Feb 2009]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US