

Product datasheet for RC201646L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IL17B Receptor (IL17RB) (NM 018725) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: IL17B Receptor (IL17RB) (NM 018725) Human Tagged ORF Clone Lentiviral Particle

Symbol: IL17B Receptor

Synonyms: CRL4; EVI27; IL17BR; IL17RH1

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_018725 **ORF Size:** 1506 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC201646).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 018725.3

RefSeq Size: 2072 bp
RefSeq ORF: 1509 bp
Locus ID: 55540
UniProt ID: Q9NRM6
Cytogenetics: 3p21.1

Protein Families: Druggable Genome, Secreted Protein, Transmembrane

Protein Pathways: Cytokine-cytokine receptor interaction





IL17B Receptor (IL17RB) (NM_018725) Human Tagged ORF Clone Lentiviral Particle – RC201646L2V

MW: 54.1 kDa

Gene Summary: The protein encoded by this gene is a cytokine receptor. This receptor specifically binds to

IL17B and IL17E, but does not bind to IL17 and IL17C. This receptor has been shown to mediate the activation of NF-kappaB and the production of IL8 induced by IL17E. The expression of the rat counterpart of this gene was found to be significantly up-regulated during intestinal inflammation, which suggested the immunoregulatory activity of this

receptor. [provided by RefSeq, Jul 2008]