

## Product datasheet for RC201660L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## KDELC1 (POGLUT2) (NM 024089) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: KDELC1 (POGLUT2) (NM\_024089) Human Tagged ORF Clone Lentiviral Particle

Symbol: POGLUT2

Synonyms: EP58; ERp58; KDEL1; KDELC1

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 024089

ORF Size: 1506 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC201660).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 024089.1, NP 076994.1

 RefSeq Size:
 2129 bp

 RefSeq ORF:
 1509 bp

 Locus ID:
 79070

 UniProt ID:
 Q6UW63

 Cytogenetics:
 13q33.1

**Domains:** Filamin, CAP10

**MW:** 58 kDa







## **Gene Summary:**

This gene encodes a protein product localized to the lumen of the endoplasmic reticulum. As a member of the endoplasmic reticulum protein family the encoded protein contains a Lys-Asp-Glu-Leu or KDEL motif located at the extreme C-terminus which prevents all endoplasmic reticulum resident proteins from being secreted. Proteins carrying this motif are bound by a receptor in the Golgi apparatus so that the receptor-ligand complex returns to the endoplasmic reticulum. A processed non-transcribed pseudogene located in an intron of a sodium transporter gene on chromosome 5 has been defined for this gene. This gene has multiple transcript variants which are predicted to encode distinct isoforms. [provided by RefSeq, Jan 2016]