

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC201678L4V

## PGAM5 (NM\_138575) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	PGAM5 (NM_138575) Human Tagged ORF Clone Lentiviral Particle
Symbol:	PGAM5
Synonyms:	BXLBV68
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_138575
ORF Size:	765 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC201678).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 138575.1</u>
RefSeq Size:	1194 bp
RefSeq ORF:	768 bp
Locus ID:	192111
UniProt ID:	<u>Q96HS1</u>
Cytogenetics:	12q24.33
Protein Families:	Transmembrane
MW:	28 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:Displays phosphatase activity for serine/threonine residues, and, dephosphorylates and<br/>activates MAP3K5 kinase. Has apparently no phosphoglycerate mutase activity. May be<br/>regulator of mitochondrial dynamics. Substrate for a KEAP1-dependent ubiquitin ligase<br/>complex. Contributes to the repression of NFE2L2-dependent gene expression. Acts as a<br/>central mediator for programmed necrosis induced by TNF, by reactive oxygen species and<br/>by calcium ionophore.[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US