

Product datasheet for RC201702L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CDC5L (NM_001253) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CDC5L (NM 001253) Human Tagged ORF Clone Lentiviral Particle

Symbol: CDC5L

Synonyms: CDC5; CDC5-LIKE; CEF1; dJ319D22.1; PCDC5RP

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_001253 **ORF Size:** 2406 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC201702).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 001253.2

 RefSeq Size:
 6432 bp

 RefSeq ORF:
 2409 bp

 Locus ID:
 988

 UniProt ID:
 Q99459

Cytogenetics: 6p21.1

Domains: myb_DNA-binding
Protein Families: Transcription Factors





CDC5L (NM_001253) Human Tagged ORF Clone Lentiviral Particle - RC201702L4V

Protein Pathways: Spliceosome

MW: 92.3 kDa

Gene Summary: The protein encoded by this gene shares a significant similarity with Schizosaccharomyces

pombe cdc5 gene product, which is a cell cycle regulator important for G2/M transition. This protein has been demonstrated to act as a positive regulator of cell cycle G2/M progression. It was also found to be an essential component of a non-snRNA spliceosome, which contains at least five additional protein factors and is required for the second catalytic step of pre-

mRNA splicing. [provided by RefSeq, Jul 2008]