

## Product datasheet for RC201797L3

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Casein Kinase 1 alpha (CSNK1A1) (NM\_001892) Human Tagged Lenti ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** Casein Kinase 1 alpha (CSNK1A1) (NM\_001892) Human Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: Casein Kinase 1 alpha

**Synonyms:** CK1; CK1a; CKIa; HEL-S-77p; HLCDGP1; PRO2975

Mammalian Cell Puromycin

Selection:

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC201797).

Sequence:

Restriction Sites: Sgfl-Mlul

**Cloning Scheme:** 





st The last codon before the Stop codon of the ORF.

**ACCN:** NM\_001892

ORF Size: 1011 bp





## Casein Kinase 1 alpha (CSNK1A1) (NM\_001892) Human Tagged Lenti ORF Clone - RC201797L3

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>NM 001892.4</u>

 RefSeq Size:
 3065 bp

 RefSeq ORF:
 1014 bp

 Locus ID:
 1452

 UniProt ID:
 P48729

Cytogenetics: 5q32

**Domains:** pkinase, TyrKc, S TKc

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Protein Kinase

**Protein Pathways:** Hedgehog signaling pathway, Wnt signaling pathway

MW: 38.7 kDa

**Gene Summary:** Casein kinases are operationally defined by their preferential utilization of acidic proteins

such as caseins as substrates. It can phosphorylate a large number of proteins. Participates in Wnt signaling. Phosphorylates CTNNB1 at 'Ser-45'. May phosphorylate PER1 and PER2. May

play a role in segregating chromosomes during mitosis (PubMed:11955436,

PubMed:1409656, PubMed:18305108). May play a role in keratin cytoskeleton disassembly and thereby, it may regulate epithelial cell migration (PubMed:23902688).[UniProtKB/Swiss-

Prot Function]