

Product datasheet for RC201805L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

FADD (NM_003824) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: FADD (NM_003824) Human Tagged ORF Clone Lentiviral Particle

Symbol: FADD

Synonyms: GIG3; MORT1

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 003824

ORF Size: 624 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC201805).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 003824.2</u>

RefSeq Size: 1855 bp
RefSeq ORF: 627 bp
Locus ID: 8772
UniProt ID: Q13158
Cytogenetics: 11q13.3

Domains: DEATH, DED

Protein Families: Druggable Genome





FADD (NM_003824) Human Tagged ORF Clone Lentiviral Particle - RC201805L1V

Protein Pathways: Alzheimer's disease, Apoptosis, Pathways in cancer, RIG-I-like receptor signaling pathway,

Toll-like receptor signaling pathway

MW: 23.3 kDa

Gene Summary: The protein encoded by this gene is an adaptor molecule that interacts with various cell

surface receptors and mediates cell apoptotic signals. Through its C-terminal death domain, this protein can be recruited by TNFRSF6/Fas-receptor, tumor necrosis factor receptor, TNFRSF25, and TNFSF10/TRAIL-receptor, and thus it participates in the death signaling initiated by these receptors. Interaction of this protein with the receptors unmasks the N-terminal effector domain of this protein, which allows it to recruit caspase-8, and thereby activate the cysteine protease cascade. Knockout studies in mice also suggest the importance

of this protein in early T cell development. [provided by RefSeq, Jul 2008]