

## Product datasheet for RC201828L1V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# SMAD4 (NM\_005359) Human Tagged ORF Clone Lentiviral Particle

#### **Product data:**

**Product Type:** Lentiviral Particles

Product Name: SMAD4 (NM 005359) Human Tagged ORF Clone Lentiviral Particle

Symbol: SMAD4

Synonyms: DPC4; JIP; MADH4; MYHRS

Mammalian Cell

Selection:

ACCN:

None

NM 005359

**Vector:** pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK

ORF Size: 1656 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC201828).

•

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 005359.3

 RefSeq Size:
 3220 bp

 RefSeq ORF:
 1659 bp

 Locus ID:
 4089

 UniProt ID:
 Q13485

Cytogenetics: 18q21.2

**Domains:** DWB, DWA, MH1

**Protein Families:** Druggable Genome, Transcription Factors





### SMAD4 (NM\_005359) Human Tagged ORF Clone Lentiviral Particle - RC201828L1V

**Protein Pathways:** Adherens junction, Cell cycle, Chronic myeloid leukemia, Colorectal cancer, Pancreatic cancer,

Pathways in cancer, TGF-beta signaling pathway, Wnt signaling pathway

**MW:** 60.3 kDa

**Gene Summary:** This gene encodes a member of the Smad family of signal transduction proteins. Smad

proteins are phosphorylated and activated by transmembrane serine-threonine receptor kinases in response to transforming growth factor (TGF)-beta signaling. The product of this gene forms homomeric complexes and heteromeric complexes with other activated Smad proteins, which then accumulate in the nucleus and regulate the transcription of target genes. This protein binds to DNA and recognizes an 8-bp palindromic sequence (GTCTAGAC) called the Smad-binding element (SBE). The protein acts as a tumor suppressor and inhibits epithelial cell proliferation. It may also have an inhibitory effect on tumors by reducing angiogenesis and increasing blood vessel hyperpermeability. The encoded protein is a crucial component of the bone morphogenetic protein signaling pathway. The Smad proteins are subject to complex regulation by post-translational modifications. Mutations or deletions in this gene have been shown to result in pancreatic cancer, juvenile polyposis syndrome, and hereditary hemorrhagic telangiectasia syndrome. [provided by RefSeq, Aug 2017]