

Product datasheet for RC201830L1

Selenophosphate synthetase 2 (SEPHS2) (NM_012248) Human Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Selenophosphate synthetase 2 (SEPHS2) (NM_012248) Human Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	Selenophosphate synthetase 2
Synonyms:	SPS2
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC201830).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.


ACCN:	NM_012248
ORF Size:	1344 bp




[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info The expression of this clone is not guaranteed due to the nature of selenoproteins.
OTI Annotation:	This clone encodes a selenoprotein containing the rare amino acid selenocysteine (Sec). Sec is encoded by UGA codon, which normally signals translational termination. Expression of this clone is not guaranteed due to the nature of selenoproteins.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_012248.2
RefSeq Size:	2351 bp
RefSeq ORF:	1347 bp
Locus ID:	22928
UniProt ID:	Q99611
Cytogenetics:	16p11.2
Domains:	AIRS
Protein Pathways:	Metabolic pathways, Selenoamino acid metabolism
Gene Summary:	This gene encodes an enzyme that catalyzes the production of monoselenophosphate (MSP) from selenide and ATP. MSP is the selenium donor required for synthesis of selenocysteine (Sec), which is co-translationally incorporated into selenoproteins at in-frame UGA codons that normally signal translation termination. The 3' UTRs of selenoprotein mRNAs contain a conserved stem-loop structure, the Sec insertion sequence (SECIS) element, which is necessary for the recognition of UGA as a Sec codon rather than as a stop signal. This protein is itself a selenoprotein containing a Sec residue at its active site, suggesting the existence of an autoregulatory mechanism. It is preferentially expressed in tissues implicated in the synthesis of selenoproteins and in sites of blood cell development. A pseudogene for this locus has been identified on chromosome 5. [provided by RefSeq, May 2017]

Product images:



Circular map for RC201830L1



Double digestion of RC201830L1 using SgfI and MluI