

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC201831L3V

PDHA1 (NM_000284) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	PDHA1 (NM_000284) Human Tagged ORF Clone Lentiviral Particle
Symbol:	PDHA1
Synonyms:	PDHA; PDHAD; PDHCE1A; PHE1A
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_000284
ORF Size:	1170 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC201831).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 000284.1</u>
RefSeq Size:	3390 bp
RefSeq ORF:	1173 bp
Locus ID:	5160
UniProt ID:	<u>P08559</u>
Cytogenetics:	Xp22.12
Domains:	E1_dehydrog
Protein Families:	Druggable Genome



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Protein Pathways:Butanoate metabolism, Citrate cycle (TCA cycle), Glycolysis / Gluconeogenesis, Metabolic
pathways, Pyruvate metabolism, Valine, leucine and isoleucine biosynthesis

43.3 kDa

MW:

Gene Summary:The pyruvate dehydrogenase (PDH) complex is a nuclear-encoded mitochondrial
multienzyme complex that catalyzes the overall conversion of pyruvate to acetyl-CoA and
CO(2), and provides the primary link between glycolysis and the tricarboxylic acid (TCA) cycle.
The PDH complex is composed of multiple copies of three enzymatic components: pyruvate
dehydrogenase (E1), dihydrolipoamide acetyltransferase (E2) and lipoamide dehydrogenase
(E3). The E1 enzyme is a heterotetramer of two alpha and two beta subunits. This gene
encodes the E1 alpha 1 subunit containing the E1 active site, and plays a key role in the
function of the PDH complex. Mutations in this gene are associated with pyruvate
dehydrogenase E1-alpha deficiency and X-linked Leigh syndrome. Alternatively spliced
transcript variants encoding different isoforms have been found for this gene.[provided by
RefSeq, Mar 2010]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US