

Product datasheet for RC201932

RAB35 (NM 006861) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: RAB35 (NM_006861) Human Tagged ORF Clone

Tag: Myc-DDK Symbol: RAB35

Synonyms: H-ray; RAB1C; RAY

Mammalian Cell N

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)ORF Nucleotide>RC201932 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

GGTGAAGCTCACGAAGAACAGTAAACGAAAGAAACGCTGCTGC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC201932 protein sequence

Red=Cloning site Green=Tags(s)

MARDYDHLFKLLIIGDSGVGKSSLLLRFADNTFSGSYITTIGVDFKIRTVEINGEKVKLQIWDTAGQERF RTITSTYYRGTHGVIVVYDVTSAESFVNVKRWLHEINONCDDVCRILVGNKNDDPERKVVETEDAYKFAG

 ${\tt QMGIQLFETSAKENVNVEEMFNCITELVLRAKKDNLAKQQQQQQNDVVKLTKNSKRKKRCC}$

TRTRPLEQKLISEEDLAANDILDYKDDDDKV



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Chromatograms: https://cdn.origene.com/chromatograms/mk6307 b06.zip

Restriction Sites:

Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM 006861

ORF Size: 603 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: NM 006861.7

RefSeq Size: 2962 bp
RefSeq ORF: 606 bp
Locus ID: 11021



UniProt ID: Q15286
Cytogenetics: 12q24.23

Protein Families: Druggable Genome


MW: 23 kDa

Gene Summary: The small GTPases Rab are key regulators of intracellular membrane trafficking, from the

formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different sets of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion. That Rab is involved in the process of endocytosis and is an essential rate-limiting regulator of the fast recycling pathway back to the plasma membrane. During cytokinesis, required for the postfurrowing terminal steps, namely for intercellular bridge stability and abscission, possibly by controlling phosphatidylinositol 4,5-bis phosphate (PIP2) and SEPT2 localization at the intercellular bridge. May indirectly regulate neurite outgrowth. Together with TBC1D13 may be involved in regulation of insulin-induced glucose transporter SLC2A4/GLUT4 translocation to the plasma membrane in adipocytes.[UniProtKB/Swiss-Prot


Function]


Product images:



Circular map for RC201932







Western blot validation of overexpression lysate (Cat# [LY416368]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC201932 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

Coomassie blue staining of purified RAB35 protein (Cat# [TP301932]). The protein was produced from HEK293T cells transfected with RAB35 cDNA clone (Cat# RC201932) using MegaTran 2.0 (Cat# [TT210002]).