

Product datasheet for RC201938L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

spindlin 1 (SPIN1) (NM 006717) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: spindlin 1 (SPIN1) (NM_006717) Human Tagged ORF Clone Lentiviral Particle

Symbol: spindlin 1

SPIN; TDRD24 Synonyms:

Mammalian Cell

Puromycin

Selection: Vector:

pLenti-C-mGFP-P2A-Puro (PS100093)

mGFP Tag:

NM 006717 ACCN:

ORF Size: 786 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC201938).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 006717.2

RefSeq Size: 4535 bp RefSeq ORF: 789 bp Locus ID: 10927 **UniProt ID:** Q9Y657 Cytogenetics: 9q22.1

Domains: Spin-Ssty MW: 29.4 kDa





Gene Summary:

Chromatin reader that specifically recognizes and binds histone H3 both trimethylated at 'Lys-4' and asymmetrically dimethylated at 'Arg-8' (H3K4me3 and H3R8me2a) and acts as an activator of Wnt signaling pathway downstream of PRMT2. In case of cancer, promotes cell cancer proliferation via activation of the Wnt signaling pathway (PubMed:24589551). Overexpression induces metaphase arrest and chromosomal instability. Localizes to active rDNA loci and promotes the expression of rRNA genes (PubMed:21960006). May play a role in cell-cycle regulation during the transition from gamete to embryo. Involved in oocyte meiotic resumption, a process that takes place before ovulation to resume meiosis of oocytes blocked in prophase I: may act by regulating maternal transcripts to control meiotic resumption.[UniProtKB/Swiss-Prot Function]