

Product datasheet for RC201973L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Fatty Acid Binding Protein 5 (FABP5) (NM_001444) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Fatty Acid Binding Protein 5 (FABP5) (NM_001444) Human Tagged ORF Clone Lentiviral

Particle

Symbol: Fatty Acid Binding Protein 5

Synonyms: E-FABP; EFABP; KFABP; PA-FABP

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_001444

ORF Size: 405 bp

ORF Nucleotide

Sequence:

Domains:

The ORF insert of this clone is exactly the same as(RC201973).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001444.1</u>

RefSeq Size: 751 bp
RefSeq ORF: 408 bp
Locus ID: 2171
UniProt ID: Q01469
Cytogenetics: 8q21.13

lipocalin





Fatty Acid Binding Protein 5 (FABP5) (NM_001444) Human Tagged ORF Clone Lentiviral Particle – RC201973L3V

Protein Pathways: PPAR signaling pathway

MW: 15.2 kDa

Gene Summary: This gene encodes the fatty acid binding protein found in epidermal cells, and was first

identified as being upregulated in psoriasis tissue. Fatty acid binding proteins are a family of small, highly conserved, cytoplasmic proteins that bind long-chain fatty acids and other hydrophobic ligands. FABPs may play roles in fatty acid uptake, transport, and metabolism. Polymorphisms in this gene are associated with type 2 diabetes. The human genome contains many pseudogenes similar to this locus.[provided by RefSeq, Feb 2011]