

Product datasheet for RC201975L3

AGA (NM_000027) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: AGA (NM_000027) Human Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: AGA

AGU; ASRG; GA Synonyms:

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

E. coli Selection: Chloramphenicol (34 ug/mL)


The ORF insert of this clone is exactly the same as(RC201975). **ORF Nucleotide**

Sequence:

Ascl-Mlul

Restriction Sites: Cloning Scheme:

> Cloning sites used for ORF Shuttling: ORF Mlu I Asc I --- GGC GCG CC C ATG --- // --- NNN ACG CGT ---



^{*} The last codon before the Stop codon of the ORF.

ACCN: NM_000027

ORF Size: 1038 bp



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



AGA (NM_000027) Human Tagged Lenti ORF Clone - RC201975L3

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 000027.2</u>, <u>NP 000018.1</u>

RefSeq Size:2113 bpRefSeq ORF:1041 bp

UniProt ID: P20933

Locus ID:

Cytogenetics:

Domains: Asparaginase 2

Protein Families: Druggable Genome, Protease

175

4q34.3

Protein Pathways: Lysosome, Other glycan degradation

MW: 37.2 kDa

Gene Summary: This gene encodes a member of the N-terminal nucleophile (Ntn) hydrolase family of

proteins. The encoded preproprotein is proteolytically processed to generate alpha and beta chains that comprise the mature enzyme. This enzyme is involved in the catabolism of N-linked oligosaccharides of glycoproteins. It cleaves asparagine from N-acetylglucosamines as one of the final steps in the lysosomal breakdown of glycoproteins. Mutations in this gene are

associated with the lysosomal storage disease aspartylglycosaminuria that results in

progressive neurodegeneration. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is subject to proteolytic processing. [provided by

RefSeq, Nov 2015]