

Product datasheet for RC201992

GRO alpha (CXCL1) (NM_001511) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: GRO alpha (CXCL1) (NM_001511) Human Tagged ORF Clone
Tag: Myc-DDK
Symbol: GRO alpha
Synonyms: FSP; GRO1; GROa; MGSA; MGSA-a; NAP-3; SCYB1
Mammalian Cell Selection: Neomycin
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
ORF Nucleotide Sequence: >RC201992 ORF sequence
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCCCGATCGCC

ATGGCCCGCGCTGCTCTCTCCGCCGCCCCAGCAATCCCCGGCTCCTGCGAGTGGCACTGCTGCTCCTGC
 TCCTGGTAGCCGCTGGCCGGCGCGCAGCAGGAGCGTCCGTGGCCACTGAACTGCGCTGCCAGTGCTTGCA
 GACCCTGCAGGGAATCACCCCAAGAACATCAAAGTGTGAACGTGAAGTCCCCGGACCCCACTGCGCC
 CAAACCGAAGTCATAGCCACACTCAAGAATGGCGGAAAGCTTGCTCAATCCTGCATCCCCATAGTTA
 AGAAAATCATCGAAAAGATGCTGAACAGTGACAAATCCAAC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC201992 protein sequence
 Red=Cloning site Green=Tags(s)
 MARAALSAAPSNPRLLRVALLLLLLVAAGRRAAGASVATELRCQCLQTLQGIHPKNIQSVNVKSPGPHCA
 QTEVIATLKNRKAACLNPA SPIVKKIIEKMLNSDKSN

TRTRPLEQKLI SEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6011_g08.zip

Restriction Sites: SgfI-MluI



[View online »](#)

Cloning Scheme:


ACCN: NM_001511

ORF Size: 321 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.


The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001511.4
RefSeq Size:	1184 bp
RefSeq ORF:	324 bp
Locus ID:	2919
UniProt ID:	P09341
Cytogenetics:	4q13.3
Domains:	IL8
Protein Families:	Druggable Genome, Secreted Protein
Protein Pathways:	Chemokine signaling pathway, Cytokine-cytokine receptor interaction, Epithelial cell signaling in Helicobacter pylori infection, NOD-like receptor signaling pathway
MW:	11.3 kDa
Gene Summary:	This antimicrobial gene encodes a member of the CXC subfamily of chemokines. The encoded protein is a secreted growth factor that signals through the G-protein coupled receptor, CXC receptor 2. This protein plays a role in inflammation and as a chemoattractant for neutrophils. Aberrant expression of this protein is associated with the growth and progression of certain tumors. A naturally occurring processed form of this protein has increased chemotactic activity. Alternate splicing results in coding and non-coding variants of this gene. A pseudogene of this gene is found on chromosome 4. [provided by RefSeq, Sep 2014]

Product images:



Circular map for RC201992