

Product datasheet for RC201998L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DBT (NM_001918) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: DBT (NM 001918) Human Tagged ORF Clone Lentiviral Particle

Symbol: DBT

Synonyms: BCATE2; BCKAD-E2; BCKADE2; BCKDH-E2; BCOADC-E2; E2B

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM_001918

 ORF Size:
 1446 bp

ORF Nucleotide

1 1 10 bp

Sequence:

The ORF insert of this clone is exactly the same as(RC201998).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 001918.2

 RefSeq Size:
 10831 bp

 RefSeq ORF:
 1449 bp

 Locus ID:
 1629

 UniProt ID:
 P11182

 Cytogenetics:
 1p21.2

Domains: biotin_lipoyl, 2-oxoacid_dh, e3_binding

Protein Families: Druggable Genome





DBT (NM_001918) Human Tagged ORF Clone Lentiviral Particle - RC201998L1V

Protein Pathways: Metabolic pathways, Valine, leucine and isoleucine degradation

MW: 53.5 kDa

Gene Summary: The branched-chain alpha-keto acid dehydrogenase complex (BCKD) is an inner-

mitochondrial enzyme complex involved in the breakdown of the branched-chain amino acids isoleucine, leucine, and valine. The BCKD complex is thought to be composed of a core of 24 transacylase (E2) subunits, and associated decarboxylase (E1), dehydrogenase (E3), and regulatory subunits. This gene encodes the transacylase (E2) subunit. Mutations in this gene result in maple syrup urine disease, type 2. Alternatively spliced transcript variants have been described, but their biological validity has not been determined. [provided by RefSeq, Jul

2008]