

Product datasheet for RC202055

TAF12 (NM 005644) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: TAF12 (NM_005644) Human Tagged ORF Clone

Tag: Myc-DDK Symbol: TAF12

Synonyms: TAF2J; TAFII20

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)ORF Nucleotide>RC202055 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGAACCAGTTTGGCCCCTCAGCCCTAATCAACCTCTCCAATTTCTCATCATAAAACCGGAACCAGCCA
GCACCCCTCCACAAGGCTCCATGGCCAATAGTACTGCAGTGGTAAAGATACCAGGCACTCCTGGGGCAGG
AGGTCGTCTTAGCCCTGAAAACAATCAGGTATTGACCAAGAAGAAATTACAGGACTTAGTAAGAGAAGTG
GATCCTAATGAGCAGTTGGATGAAGATGTGGAGGAGATGCTGCTGCAGATTGCTGATGATTTTATCGAGA
GTGTGGTGACAGCCAGCCTGTCAGCTTGCGCGGCATCGCAAGTCTAGCACCCTGGAGGTGAAAGATGTCCA
GCTGCATTTAGAGCGCCCAGTGGAACATGTGGATCCCAGGATTTGGCTCTGAAGAAATCCGACCCTACAAA
AAAGCTTGCACCACAGAAGCTCACAAACAGAGAATGGCATTGATCCGGAAAACAACCAAGAAA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC202055 protein sequence

Red=Cloning site Green=Tags(s)

MNQFGPSALINLSNFSSIKPEPASTPPQGSMANSTAVVKIPGTPGAGGRLSPENNQVLTKKKLQDLVREV DPNEQLDEDVEEMLLQIADDFIESVVTAACQLARHRKSSTLEVKDVQLHLERQWNMWIPGFGSEEIRPYK

KACTTEAHKQRMALIRKTTKK

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6304_e12.zip



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM 005644

ORF Size: 483 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 005644.4</u>

RefSeq Size: 1129 bp RefSeq ORF: 486 bp Locus ID: 6883 UniProt ID: Q16514

Cytogenetics: 1p35.3

Domains: TFIID_A

Protein Families: Transcription Factors


Protein Pathways: Basal transcription factors

MW: 17.9 kDa

Gene Summary: Control of transcription by RNA polymerase II involves the basal transcription machinery


which is a collection of proteins. These proteins with RNA polymerase II, assemble into complexes which are modulated by transactivator proteins that bind to cis-regulatory elements located adjacent to the transcription start site. Some modulators interact directly with the basal complex, whereas others may act as bridging proteins linking transactivators to the basal transcription factors. Some of these associated factors are weakly attached while others are tightly associated with TBP in the TFIID complex. Among the latter are the TAF proteins. Different TAFs are predicted to mediate the function of distinct transcriptional activators for a variety of gene promoters and RNA polymerases. TAF12 interacts directly with TBP as well as with TAF2I. Two transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Sep 2008]


Product images:



Circular map for RC202055







Western blot validation of overexpression lysate (Cat# [LY427599]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC225149] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

Coomassie blue staining of purified TAF12 protein (Cat# [TP302055]). The protein was produced from HEK293T cells transfected with TAF12 cDNA clone (Cat# RC202055) using MegaTran 2.0 (Cat# [TT210002]).