

Product datasheet for RC202059L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Caspase-7 (CASP7) (NM_033339) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Caspase-7 (CASP7) (NM_033339) Human Tagged ORF Clone Lentiviral Particle

Symbol: Caspase-7

Synonyms: CASP-7; CMH-1; ICE-LAP3; LICE2; MCH3

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK

ACCN: NM_033339

ORF Size: 909 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC202059).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 033339.3

 RefSeq Size:
 2681 bp

 RefSeq ORF:
 912 bp

 Locus ID:
 840

 UniProt ID:
 P55210

Cytogenetics: 10q25.3

Protein Families: Druggable Genome, Protease

Protein Pathways: Alzheimer's disease, Apoptosis





MW: 34.3 kDa

Gene Summary: This gene encodes a member of the cysteine-aspartic acid protease (caspase) family.

Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. The precursor of the encoded protein is cleaved by caspase 3 and 10, is activated upon cell death stimuli and induces apoptosis. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, May 2012]