

Product datasheet for RC202061

ID1 (NM_002165) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: ID1 (NM_002165) Human Tagged ORF Clone

Tag: Myc-DDK

Symbol: ID1

Synonyms: bHLHb24; ID

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)ORF Nucleotide>RC202061 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC202061 protein sequence

Red=Cloning site Green=Tags(s)

MKVASGSTATAAAGPSCALKAGKTASGAGEVVRCLSEQSVAISRCAGGAGARLPALLDEQQVNVLLYDMN GCYSRLKELVPTLPQNRKVSKVEILQHVIDYIRDLQLELNSESEVGTPGGRGLPVRAPLSTLNGEISALT

AEAACVPADDRILCR

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6011 a04.zip



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com ORIGENE

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_002165

ORF Size: 465 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts

of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at customercom or by

calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 002165.4</u>

 RefSeq Size:
 1000 bp

 RefSeq ORF:
 468 bp

 Locus ID:
 3397

 UniProt ID:
 P41134

 Cytogenetics:
 20q11.21

Domains: HLH

Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: TGF-beta signaling pathway

MW: 16.1 kDa

Gene Summary: The protein encoded by this gene is a helix-loop-helix (HLH) protein that can form


heterodimers with members of the basic HLH family of transcription factors. The encoded

protein has no DNA binding activity and therefore can inhibit the DNA binding and


transcriptional activation ability of basic HLH proteins with which it interacts. This protein may play a role in cell growth, senescence, and differentiation. Two transcript variants encoding


different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

Product images:



Circular map for RC202061





Western blot validation of overexpression lysate (Cat# [LY400785]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC202061 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

Coomassie blue staining of purified ID1 protein (Cat# [TP302061]). The protein was produced from HEK293T cells transfected with ID1 cDNA clone (Cat# RC202061) using MegaTran 2.0 (Cat# [TT210002]).