

## Product datasheet for **RC202079**

### IL1 beta (IL1B) (NM\_000576) Human Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	IL1 beta (IL1B) (NM_000576) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	IL1B
Synonyms:	IL-1; IL1-BETA; IL1beta; IL1F2
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RC202079 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGGCAGAAGTACCTGAGCTCGCCAGTGAAATGATGGCTTATTACAGTGGCAATGAGGATGACTTGTCT  
TTGAAGCTGATGGCCCTAACAGATGAAGTCTCCTTCCAGGACCTGGACCTCTGCCCTCTGGATGGCGG  
CATCCAGCTACGAATCTCCGACCACCACTACAGCAAGGGCTTCAGGCAGGCCGCGTCAGTTGTTGTGGCC  
ATGGACAAGCTGAGGAAGATGCTGGTCCCTGCCACAGACCTTCCAGGAGAATGACCTGAGCACCTTCT  
TTCCCTTCATCTTGAAGAAGAACCTATCTTCTCGACACATGGGATAACGAGGCTTATGTGCACGATGC  
ACCTGTACGATCACTGAAGTGCACGCTCCGGGACTCACAGCAAAAAGCTTGGTGATGTCTGGTCCATAT  
GAACTGAAAGCTCTCCACCTCCAGGGACAGGATATGGAGCAACAAGTGGTGTCTCCATGTCTTTGTAC  
AAGGAGAAGAAAGTAATGACAAAATACCTGTGGCCTTGGGCTCAAGGAAAAGAATCTGTACCTGTCTGT  
CGTGTGAAAGATGATAAGCCACTCTACAGCTGGAGAGTGTAGATCCCAAAAATACCCAAAGAAGAAG  
ATGGAAAAGCGATTTGTCTTCAACAAGATAGAAATCAATAACAAGCTGGAATTTGAGTCTGCCAGTTCC  
CCAAGTGGTACATCAGCACCTCTAAGCAGAAAACATGCCGCTTCTCTGGGAGGACCAAAAGCGGCCA  
GGATATAACTGACTTCACCATGCAATTTGTGTCTTCC

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

**Protein Sequence:** >RC202079 protein sequence  
Red=Cloning site Green=Tags(s)

MAEVP ELASEMMAYYS GNEDDLFFEADGPKQMKCSFQDL DLCPLDGGIQLRISDHHYSKGRQAASVVVA  
 MDKLRKMLVPCPQTFQENDLSTFFPFI FEEPIFFDTWDNEAYVHDAPVRS LNCTLRDSQQKSLVMSGPY  
 ELKALHLQGQDMEQQVVF SMSFVQGEESNDKIPVALGLKEKNL YLSCVLKDDKPTLQLESVDPKNYPKKK  
 MEKRFVFNKIEINNKL EFESAQFPNWIISTSAENMPVFLGGTKGGQDITDFTMQFVSS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


**Chromatograms:** [https://cdn.origene.com/chromatograms/mk6125\\_d05.zip](https://cdn.origene.com/chromatograms/mk6125_d05.zip)

**Restriction Sites:** SgfI-MluI

**Cloning Scheme:**



**Plasmid Map:**




**ACCN:** NM\_000576

**ORF Size:** 807 bp


<b>OTI Disclaimer:</b>	<p>Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:custsupport@origene.com">custsupport@origene.com</a> or by calling 301.340.3188 option 3 for pricing and delivery.</p> <p>The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a></p>
<b>OTI Annotation:</b>	<p>This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.</p>
<b>Components:</b>	<p>The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).</p>
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_000576.3</a>
<b>RefSeq Size:</b>	1498 bp
<b>RefSeq ORF:</b>	810 bp
<b>Locus ID:</b>	3553
<b>UniProt ID:</b>	<a href="#">P01584</a>
<b>Cytogenetics:</b>	2q14.1
<b>Protein Families:</b>	Druggable Genome, Secreted Protein
<b>Protein Pathways:</b>	Alzheimer's disease, Apoptosis, Cytokine-cytokine receptor interaction, Cytosolic DNA-sensing pathway, Graft-versus-host disease, Hematopoietic cell lineage, MAPK signaling pathway, NOD-like receptor signaling pathway, Prion diseases, Toll-like receptor signaling pathway, Type I diabetes mellitus
<b>MW:</b>	30.7 kDa

**Gene Summary:**


The protein encoded by this gene is a member of the interleukin 1 cytokine family. This cytokine is produced by activated macrophages as a proprotein, which is proteolytically processed to its active form by caspase 1 (CASP1/ICE). This cytokine is an important mediator of the inflammatory response, and is involved in a variety of cellular activities, including cell proliferation, differentiation, and apoptosis. The induction of cyclooxygenase-2 (PTGS2/COX2) by this cytokine in the central nervous system (CNS) is found to contribute to inflammatory pain hypersensitivity. Similarly, IL-1B has been implicated in human osteoarthritis pathogenesis. Patients with severe Coronavirus Disease 2019 (COVID-19) present elevated levels of pro-inflammatory cytokines such as IL-1B in bronchial alveolar lavage fluid samples. The lung damage induced by the Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is to a large extent, a result of the inflammatory response promoted by cytokines such as IL-1B. This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2. [provided by RefSeq, Jul 2020]

**Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY IL1B (Cat# RC202079, Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IL1B (Cat# [TA506443]). Positive lysates [LY400196] (100ug) and [LC400196] (20ug) can be purchased separately from OriGene.



Western blot validation of overexpression lysate (Cat# [LY400196]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC202079 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).



Coomassie blue staining of purified IL1B protein (Cat# [TP302079]). The protein was produced from HEK293T cells transfected with IL1B cDNA clone (Cat# RC202079) using MegaTran 2.0 (Cat# [TT210002]).



Anti-IL1B mouse monoclonal antibody ([TA506443]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY IL1B (RC202079).