

Product datasheet for RC202105L4

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

BPGM (NM_199186) Human Tagged Lenti ORF Clone

Puromycin

Sgfl-Mlul

Product data:

Product Type: Expression Plasmids

Product Name: BPGM (NM_199186) Human Tagged Lenti ORF Clone

Tag: mGFP Symbol: BPGM

Synonyms: DPGM; ECYT8

Mammalian Cell

Selection:

Vector:

pLenti-C-mGFP-P2A-Puro (PS100093)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC202105).


Sequence:

Restriction Sites: Cloning Scheme:

Cloning sites used for ORF Shuttling:

Sgf I ORF MIu I

GGG ATC GC ATG ---//--- NNÑ ACG CGT ---



^{*} The last codon before the Stop codon of the ORF.

ACCN: NM_199186

ORF Size: 777 bp



BPGM (NM_199186) Human Tagged Lenti ORF Clone - RC202105L4

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 199186.1</u>, <u>NP 954655.1</u>

RefSeq Size: 2121 bp
RefSeq ORF: 780 bp
Locus ID: 669

 UniProt ID:
 P07738

 Cytogenetics:
 7q33

Protein Families: Druggable Genome

Protein Pathways: Glycolysis / Gluconeogenesis, Metabolic pathways

MW: 30 kDa


Gene Summary: 2,3-diphosphoglycerate (2,3-DPG) is a small molecule found at high concentrations in red

blood cells where it binds to and decreases the oxygen affinity of hemoglobin. This gene encodes a multifunctional enzyme that catalyzes 2,3-DPG synthesis via its synthetase activity, and 2,3-DPG degradation via its phosphatase activity. The enzyme also has phosphoglycerate phosphomutase activity. Deficiency of this enzyme increases the affinity of cells for oxygen. Mutations in this gene result in hemolytic anemia. Multiple alternatively spliced variants,

encoding the same protein, have been identified. [provided by RefSeq, Sep 2009]



Product images:



Circular map for RC202105L4