

Product datasheet for RC202143L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MTHFD2 (NM 006636) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: MTHFD2 (NM_006636) Human Tagged ORF Clone Lentiviral Particle

Symbol: **NMDMC** Synonyms:

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Myc-DDK Tag: NM 006636 ACCN: **ORF Size:** 1050 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC202143).

The molecular sequence of this clone aligns with the gene accession number as a point of OTI Disclaimer: reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 006636.3

RefSeq Size: 2208 bp RefSeq ORF: 1053 bp Locus ID: 10797 **UniProt ID:** P13995 Cytogenetics: 2p13.1

Domains: THF_DHG_CYH

Protein Families: Druggable Genome





MTHFD2 (NM_006636) Human Tagged ORF Clone Lentiviral Particle - RC202143L3V

Protein Pathways: Glyoxylate and dicarboxylate metabolism, Metabolic pathways, One carbon pool by folate

MW: 37.9 kDa

Gene Summary: This gene encodes a nuclear-encoded mitochondrial bifunctional enzyme with

methylenetetrahydrofolate dehydrogenase and methenyltetrahydrofolate cyclohydrolase activities. The enzyme functions as a homodimer and is unique in its absolute requirement for magnesium and inorganic phosphate. Formation of the enzyme-magnesium complex allows binding of NAD. Alternative splicing results in two different transcripts, one protein-coding and the other not protein-coding. This gene has a pseudogene on chromosome 7.

[provided by RefSeq, Mar 2009]