

Product datasheet for **RC202180L2V**

MCP1 (CCL2) (NM_002982) Human Tagged ORF Clone Lentiviral Particle

Product data:

| | |
|---------------------------|--|
| Product Type: | Lentiviral Particles |
| Product Name: | MCP1 (CCL2) (NM_002982) Human Tagged ORF Clone Lentiviral Particle |
| Symbol: | MCP1 |
| Synonyms: | GDCE-2; HC11; HSMCR30; MCAF; MCP-1; MCP1; SCYA2; SMC-CF |
| Mammalian Cell Selection: | None |
| Vector: | pLenti-C-mGFP (PS100071) |
| Tag: | mGFP |
| ACCN: | NM_002982 |
| ORF Size: | 297 bp |
| ORF Nucleotide Sequence: | The ORF insert of this clone is exactly the same as(RC202180). |
| OTI Disclaimer: | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info |
| OTI Annotation: | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. |
| RefSeq: | NM_002982.3 |
| RefSeq Size: | 760 bp |
| RefSeq ORF: | 300 bp |
| Locus ID: | 6347 |
| UniProt ID: | P13500 |
| Cytogenetics: | 17q12 |
| Domains: | IL8 |
| Protein Families: | Druggable Genome, Secreted Protein |



[View online »](#)

| | |
|--------------------------|--|
| Protein Pathways: | Chemokine signaling pathway, Cytokine-cytokine receptor interaction, NOD-like receptor signaling pathway |
| MW: | 11 kDa |
| Gene Summary: | <p>This gene is one of several cytokine genes clustered on the q-arm of chromosome 17. Chemokines are a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of N-terminal cysteine residues of the mature peptide. This chemokine is a member of the CC subfamily which is characterized by two adjacent cysteine residues. This cytokine displays chemotactic activity for monocytes and basophils but not for neutrophils or eosinophils. It has been implicated in the pathogenesis of diseases characterized by monocytic infiltrates, like psoriasis, rheumatoid arthritis and atherosclerosis. It binds to chemokine receptors CCR2 and CCR4. Elevated expression of the encoded protein is associated with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection. [provided by RefSeq, Aug 2020]</p> |