

## Product datasheet for RC202182L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## IKB beta (NFKBIB) (NM\_002503) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: IKB beta (NFKBIB) (NM\_002503) Human Tagged ORF Clone Lentiviral Particle

Symbol: IKB beta
Synonyms: IKBB; TRIP9

Mammalian Cell F

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_002503 **ORF Size:** 1068 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC202182).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 002503.3

 RefSeq Size:
 1243 bp

 RefSeq ORF:
 1071 bp

 Locus ID:
 4793

 UniProt ID:
 Q15653

Cytogenetics: 19q13.2

Domains: ANK

**Protein Families:** Stem cell - Pluripotency, Transcription Factors





## IKB beta (NFKBIB) (NM\_002503) Human Tagged ORF Clone Lentiviral Particle - RC202182L4V

**Protein Pathways:** Adipocytokine signaling pathway, B cell receptor signaling pathway, Chemokine signaling

pathway, Cytosolic DNA-sensing pathway, Neurotrophin signaling pathway, NOD-like receptor signaling pathway, RIG-I-like receptor signaling pathway, T cell receptor signaling pathway

MW: 37.8 kDa

**Gene Summary:** The protein encoded by this gene belongs to the NF-kappa-B inhibitor family, which inhibit

NF-kappa-B by complexing with, and trapping it in the cytoplasm. Phosphorylation of serine residues on these proteins by kinases marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NF-kappa-B, which translocates to the nucleus to function as a transcription factor. Alternatively spliced transcript variants have been found

for this gene.[provided by RefSeq, Jul 2011]