

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC202195L4V

## Cystathionase (CTH) (NM\_001902) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	Cystathionase (CTH) (NM_001902) Human Tagged ORF Clone Lentiviral Particle
Symbol:	Cystathionase
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_001902
ORF Size:	1215 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC202195).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001902.4</u>
RefSeq Size:	2140 bp
RefSeq ORF:	1218 bp
Locus ID:	1491
UniProt ID:	<u>P32929</u>
Cytogenetics:	1p31.1
Protein Pathways:	Cysteine and methionine metabolism, Glycine, serine and threonine metabolism, Metabolic pathways, Nitrogen metabolism, Selenoamino acid metabolism
MW:	44.5 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:This gene encodes a cytoplasmic enzyme in the trans-sulfuration pathway that converts<br/>cystathione derived from methionine into cysteine. Glutathione synthesis in the liver is<br/>dependent upon the availability of cysteine. Mutations in this gene cause cystathioninuria.<br/>Alternative splicing of this gene results in three transcript variants encoding different<br/>isoforms. [provided by RefSeq, Jun 2010]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US