

Product datasheet for RC202245L1

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

COX2 (PTGS2) (NM_000963) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: COX2 (PTGS2) (NM_000963) Human Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: COX2

Synonyms: COX-2; COX2; GRIPGHS; hCox-2; PGG/HS; PGHS-2; PHS-2

Mammalian Cell None

Selection:

Vector:pLenti-C-Myc-DDK (PS100064)E. coli Selection:Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC202245).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_000963

ORF Size: 1812 bp



COX2 (PTGS2) (NM_000963) Human Tagged Lenti ORF Clone - RC202245L1

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 000963.1</u>

 RefSeq Size:
 4465 bp

 RefSeq ORF:
 1815 bp

 Locus ID:
 5743

 UniProt ID:
 P35354

Cytogenetics: 1q31.1

Domains: An_peroxidase, EGF
Protein Families: Druggable Genome

Protein Pathways: Arachidonic acid metabolism, Pathways in cancer, Small cell lung cancer, VEGF signaling

pathway

MW: 69 kDa

Gene Summary: Prostaglandin-endoperoxide synthase (PTGS), also known as cyclooxygenase, is the key

enzyme in prostaglandin biosynthesis, and acts both as a dioxygenase and as a peroxidase. There are two isozymes of PTGS: a constitutive PTGS1 and an inducible PTGS2, which differ in


their regulation of expression and tissue distribution. This gene encodes the inducible

isozyme. It is regulated by specific stimulatory events, suggesting that it is responsible for the prostanoid biosynthesis involved in inflammation and mitogenesis. [provided by RefSeq, Feb


20091



Product images:



Circular map for RC202245L1



Double digestion of RC202245L1 using Sgfl and Mlul