

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC202245L4V

COX2 (PTGS2) (NM_000963) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	COX2 (PTGS2) (NM_000963) Human Tagged ORF Clone Lentiviral Particle
Symbol:	PTGS2
Synonyms:	COX-2; COX2; GRIPGHS; hCox-2; PGG/HS; PGHS-2; PHS-2
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_000963
ORF Size:	1812 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC202245).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 000963.1</u>
RefSeq Size:	4465 bp
RefSeq ORF:	1815 bp
Locus ID:	5743
UniProt ID:	<u>P35354</u>
Cytogenetics:	1q31.1
Domains:	An_peroxidase, EGF
Protein Families:	Druggable Genome



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE COX2 (PTGS2) (NM_000963) Human Tagged ORF Clone Lentiviral Particle – RC202245L4V	
Protein Pathways:	Arachidonic acid metabolism, Pathways in cancer, Small cell lung cancer, VEGF signaling pathway
MW:	69 kDa
Gene Summary:	Prostaglandin-endoperoxide synthase (PTGS), also known as cyclooxygenase, is the key enzyme in prostaglandin biosynthesis, and acts both as a dioxygenase and as a peroxidase. There are two isozymes of PTGS: a constitutive PTGS1 and an inducible PTGS2, which differ in their regulation of expression and tissue distribution. This gene encodes the inducible isozyme. It is regulated by specific stimulatory events, suggesting that it is responsible for the prostanoid biosynthesis involved in inflammation and mitogenesis. [provided by RefSeq, Feb 2009]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US