

Product datasheet for **RC202282**

IL15 (NM_000585) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: IL15 (NM_000585) Human Tagged ORF Clone
Tag: Myc-DDK
Symbol: IL15
Synonyms: IL-15
Mammalian Cell Selection: Neomycin
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
ORF Nucleotide Sequence: >RC202282 ORF sequence
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGAGAATTTGAAACCACATTTGAGAAGTATTTCCATCCAGTGCTACTTGTGTTTACTTCTAAACAGTC
ATTTTCTAACTGAAGCTGGCATTGCTCTTCAATTTGGGCTGTTTCAGTGCAGGGCTTCTAAAACAGA
AGCCAACGGGTGAATGTAATAAGTGATTTGAAAAAATTGAAGATCTTATTCAATCTATGCATATTGAT
GCTACTTTATATACGAAAGTGATGTTACCCAGTTGCAAAGTAACAGCAATGAAGTGCTTTCTCTTGG
AGTTACAAGTTATTTCACTTGAGTCCGGAGATGCAAGTATTCATGATACAGTAGAAAATCTGATCATCCT
AGCAAACAACAGTTTGTCTTCTAATGGGAATGTAAACAGAATCTGGATGCAAAGAATGTGAGGAACCTGGAG
GAAAAAATATTAAGAATTTTTGCAGAGTTTTGTACATATTGCCAAATGTTTCATCAACACTTCT

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC202282 protein sequence
Red=Cloning site Green=Tags(s)

MRISKPHLRISIQCYLCLLNHFLTEAGIHVFIILGCF SAGLPKTEANWVVISDLKKIEDLIQSMHID
ATLYTESDVHPSCKVTAMKCFLELQVISLESGDASIHDVTENLIILANNSLSSNGNVTESGCKECELE
EKNIKEFLQSFVHIVQMFINTS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6523_b06.zip



[View online >](#)

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_000585

ORF Size: 486 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.


The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_000585.5
RefSeq Size:	2012 bp
RefSeq ORF:	489 bp
Locus ID:	3600
UniProt ID:	P40933
Cytogenetics:	4q31.21
Protein Families:	Druggable Genome, Secreted Protein
Protein Pathways:	Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway
MW:	18.1 kDa
Gene Summary:	<p>The protein encoded by this gene is a cytokine that regulates T and natural killer cell activation and proliferation. This cytokine and interleukine 2 share many biological activities. They are found to bind common hematopoietin receptor subunits, and may compete for the same receptor, and thus negatively regulate each other's activity. The number of CD8+ memory cells is shown to be controlled by a balance between this cytokine and IL2. This cytokine induces the activation of JAK kinases, as well as the phosphorylation and activation of transcription activators STAT3, STAT5, and STAT6. Studies of the mouse counterpart suggested that this cytokine may increase the expression of apoptosis inhibitor BCL2L1/BCL-x(L), possibly through the transcription activation activity of STAT6, and thus prevent apoptosis. Alternatively spliced transcript variants of this gene have been reported. [provided by RefSeq, Feb 2011]</p>

Product images:



Circular map for RC202282