

Product datasheet for RC202318L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

KCNMB4 (NM_014505) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: KCNMB4 (NM_014505) Human Tagged ORF Clone Lentiviral Particle

Symbol: KCNMB4

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_014505

ORF Size: 630 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC202318).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 014505.4</u>

 RefSeq Size:
 4725 bp

 RefSeq ORF:
 633 bp

 Locus ID:
 27345

 UniProt ID:
 Q86W47

 Cytogenetics:
 12q15

 Domains:
 CaKB

Protein Families: Druggable Genome, Ion Channels: Other, Transmembrane

Protein Pathways: Vascular smooth muscle contraction





KCNMB4 (NM_014505) Human Tagged ORF Clone Lentiviral Particle - RC202318L4V

MW: 23.9 kDa

Gene Summary: MaxiK channels are large conductance, voltage and calcium-sensitive potassium channels

which are fundamental to the control of smooth muscle tone and neuronal excitability. MaxiK channels can be formed by 2 subunits: the pore-forming alpha subunit and the modulatory beta subunit. The protein encoded by this gene is an auxiliary beta subunit which slows activation kinetics, leads to steeper calcium sensitivity, and shifts the voltage range of current activation to more negative potentials than does the beta 1 subunit. [provided by RefSeq, Jul

2008]