

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC202332L4V

## MXI1 (NM\_001008541) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	MXI1 (NM_001008541) Human Tagged ORF Clone Lentiviral Particle
Symbol:	MXI1
Synonyms:	bHLHc11; MAD2; MXD2; MXI
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_001008541
ORF Size:	546 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC202332).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001008541.1, NP 001008541.1</u>
RefSeq Size:	3047 bp
RefSeq ORF:	549 bp
Locus ID:	4601
UniProt ID:	<u>P50539</u>
Cytogenetics:	10q25.2
Protein Families:	Druggable Genome, Transcription Factors
MW:	20.6 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:Expression of the c-myc gene, which produces an oncogenic transcription factor, is tightly<br/>regulated in normal cells but is frequently deregulated in human cancers. The protein<br/>encoded by this gene is a transcriptional repressor thought to negatively regulate MYC<br/>function, and is therefore a potential tumor suppressor. This protein inhibits the<br/>transcriptional activity of MYC by competing for MAX, another basic helix-loop-helix protein<br/>that binds to MYC and is required for its function. Defects in this gene are frequently found in<br/>patients with prostate tumors. Three alternatively spliced transcripts encoding different<br/>isoforms have been described. Additional alternatively spliced transcripts may exist but the<br/>products of these transcripts have not been verified experimentally. [provided by RefSeq, Jul<br/>2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US