

## Product datasheet for **RC202340L1V**

### APC16 (ANAPC16) (NM\_173473) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

|                           |  |
|---------------------------|--|
| Product Type:             | Lentiviral Particles   |
| Product Name:             | APC16 (ANAPC16) (NM_173473) Human Tagged ORF Clone Lentiviral Particle   |
| Symbol:                   | APC16  |
| Synonyms:                 | APC16; bA570G20.3; C10orf104; CENP-27; MSAG  |
| Mammalian Cell Selection: | None   |
| Vector:                   | pLenti-C-Myc-DDK (PS100064)  |
| Tag:                      | Myc-DDK  |
| ACCN:                     | NM_173473  |
| ORF Size:                 | 330 bp   |
| ORF Nucleotide Sequence:  | The ORF insert of this clone is exactly the same as(RC202340).   |
| OTI Disclaimer:           | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a> |
| OTI Annotation:           | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.   |
| RefSeq:                   | <a href="#">NM_173473.2</a>  |
| RefSeq Size:              | 3252 bp  |
| RefSeq ORF:               | 333 bp   |
| Locus ID:                 | 119504   |
| UniProt ID:               | <a href="#">Q96DE5</a>   |
| Cytogenetics:             | 10q22.1  |
| MW:                       | 11.7 kDa   |



[View online »](#)

**Gene Summary:**

Component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle. The APC/C complex acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains. [UniProtKB/Swiss-Prot Function]