

## Product datasheet for RC202368L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## RFX5 (NM\_000449) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

**Product Type: Lentiviral Particles** 

**Product Name:** RFX5 (NM\_000449) Human Tagged ORF Clone Lentiviral Particle

Symbol: RFX5

**Mammalian Cell** Puromycin

Selection:

Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM\_000449

**ORF Size:** 1848 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC202368).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 000449.3, NP 000440.1

RefSeq Size: 3618 bp RefSeq ORF: 1851 bp Locus ID: 5993 **UniProt ID:** P48382 **Cytogenetics:** 1q21.3

Domains: RFX\_DNA\_binding **Protein Families: Transcription Factors** 

**Protein Pathways:** Antigen processing and presentation, Primary immunodeficiency







**MW:** 65.1 kDa

**Gene Summary:** 

A lack of MHC-II expression results in a severe immunodeficiency syndrome called MHC-II deficiency, or the bare lymphocyte syndrome (BLS; MIM 209920). At least 4 complementation groups have been identified in B-cell lines established from patients with BLS. The molecular defects in complementation groups B, C, and D all lead to a deficiency in RFX, a nuclear protein complex that binds to the X box of MHC-II promoters. The lack of RFX binding activity in complementation group C results from mutations in the RFX5 gene encoding the 75-kD subunit of RFX (Steimle et al., 1995). RFX5 is the fifth member of the growing family of DNA-binding proteins sharing a novel and highly characteristic DNA-binding domain called the RFX motif. Multiple alternatively spliced transcript variants have been found but the full-length natures of only two have been determined. [provided by RefSeq, Jul 2008]