

Product datasheet for RC202390L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IL17RA (NM_014339) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: IL17RA (NM_014339) Human Tagged ORF Clone Lentiviral Particle

Symbol: IL17RA

Synonyms: CANDF5; CD217; CDw217; hlL-17R; IL-17RA; IL17R; IMD51

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 014339

ORF Size: 2598 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC202390).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 014339.4

 RefSeq Size:
 3429 bp

 RefSeq ORF:
 2601 bp

 Locus ID:
 23765

 UniProt ID:
 Q96F46

 Cytogenetics:
 22q11.1

Protein Families: Druggable Genome, Transmembrane
Protein Pathways: Cytokine-cytokine receptor interaction





ORIGENE

MW: 96.13 kDa

Gene Summary: Interleukin 17A (IL17A) is a proinflammatory cytokine secreted by activated T-lymphocytes. It

is a potent inducer of the maturation of CD34-positive hematopoietic precursors into neutrophils. The transmembrane protein encoded by this gene (interleukin 17A receptor; IL17RA) is a ubiquitous type I membrane glycoprotein that binds with low affinity to

interleukin 17A. Interleukin 17A and its receptor play a pathogenic role in many inflammatory and autoimmune diseases such as rheumatoid arthritis. Like other cytokine receptors, this receptor likely has a multimeric structure. Alternative splicing results in multiple transcript

variants encoding different isoforms. [provided by RefSeq, Feb 2014]