

## Product datasheet for **RC202437L2V**

### Cyclin D3 (CCND3) (NM\_001760) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	Cyclin D3 (CCND3) (NM_001760) Human Tagged ORF Clone Lentiviral Particle
Symbol:	Cyclin D3
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
Tag:	mGFP
ACCN:	NM_001760
ORF Size:	876 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC202437).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_001760.2</a>
RefSeq Size:	2095 bp
RefSeq ORF:	879 bp
Locus ID:	896
UniProt ID:	<a href="#">P30281</a>
Cytogenetics:	6p21.1
Domains:	cyclin_C, CYCLIN, cyclin
Protein Families:	Druggable Genome



[View online »](#)

<b>Protein Pathways:</b>	Cell cycle, Focal adhesion, Jak-STAT signaling pathway, p53 signaling pathway, Wnt signaling pathway
<b>MW:</b>	32.3 kDa
<b>Gene Summary:</b>	<p>The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK4 or CDK6, whose activity is required for cell cycle G1/S transition. This protein has been shown to interact with and be involved in the phosphorylation of tumor suppressor protein Rb. The CDK4 activity associated with this cyclin was reported to be necessary for cell cycle progression through G2 phase into mitosis after UV radiation. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2008]</p>