

Product datasheet for RC202459L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Chemokine Receptor D6 (ACKR2) (NM_001296) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Chemokine Receptor D6 (ACKR2) (NM 001296) Human Tagged ORF Clone Lentiviral Particle

Symbol: Chemokine Receptor D6

Synonyms: CCBP2; CCR9; CCR10; CMKBR9; D6; hD6

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_001296

ORF Size: 1152 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC202459).

OTI Disclaimer:

Sequence:

Domains:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 001296.3

 RefSeq Size:
 2978 bp

 RefSeq ORF:
 1155 bp

 Locus ID:
 1238

 UniProt ID:
 000590

 Cytogenetics:
 3p22.1

Protein Families: Druggable Genome, GPCR, Transmembrane

7tm 1





Chemokine Receptor D6 (ACKR2) (NM_001296) Human Tagged ORF Clone Lentiviral Particle – RC202459L3V

MW: 43.4 kDa

Gene Summary:

This gene encodes a beta chemokine receptor, which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. Chemokines and their receptor-mediated signal transduction are critical for the recruitment of effector immune cells to the inflammation site. This gene is expressed in a range of tissues and hemopoietic cells. The expression of this receptor in lymphatic endothelial cells and overexpression in vascular tumors suggested its function in chemokine-driven recirculation of leukocytes and possible chemokine effects on the development and growth of vascular tumors. This receptor appears to bind the majority of beta-chemokine family members; however, its specific function remains unknown. This gene is mapped to chromosome 3p21.3, a region that includes a cluster of chemokine receptor genes. [provided by RefSeq, Jul 2008]