

Product datasheet for RC202461L2

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

HIF-1 alpha (HIF1A) (NM_001530) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: HIF-1 alpha (HIF1A) (NM_001530) Human Tagged Lenti ORF Clone

Tag: mGFP

Symbol: HIF-1 alpha

Synonyms: bHLHe78; HIF-1-alpha; HIF-1A; HIF-1alpha; HIF1; HIF1-ALPHA; MOP1; PASD8

Mammalian Cell None

Selection:

Vector: pLenti-C-mGFP (PS100071)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC202461).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





st The last codon before the Stop codon of the ORF.

ACCN: NM_001530

ORF Size: 2478 bp



HIF-1 alpha (HIF1A) (NM_001530) Human Tagged Lenti ORF Clone - RC202461L2

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001530.2</u>

 RefSeq Size:
 3958 bp

 RefSeq ORF:
 2481 bp

 Locus ID:
 3091

 UniProt ID:
 016665

Cytogenetics: <u>Q16665</u>

Domains: PAS, HLH, PAC

Protein Families: Transcription Factors

Protein Pathways: mTOR signaling pathway, Pathways in cancer, Renal cell carcinoma

MW: 92.5 kDa

Gene Summary: This gene encodes the alpha subunit of transcription factor hypoxia-inducible factor-1 (HIF-1),

which is a heterodimer composed of an alpha and a beta subunit. HIF-1 functions as a master regulator of cellular and systemic homeostatic response to hypoxia by activating transcription of many genes, including those involved in energy metabolism, angiogenesis, apoptosis, and other genes whose protein products increase oxygen delivery or facilitate


metabolic adaptation to hypoxia. HIF-1 thus plays an essential role in embryonic

vascularization, tumor angiogenesis and pathophysiology of ischemic disease. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene.


[provided by RefSeq, Jul 2011]



Product images:



Circular map for RC202461L2



Double digestion of RC202461L2 using Sgfl and Mlul