

## Product datasheet for RC202461L3

### HIF-1 alpha (HIF1A) (NM\_001530) Human Tagged Lenti ORF Clone

#### Product data:

|                           |   |
|---------------------------|---|
| Product Type:             | Expression Plasmids   |
| Product Name:             | HIF-1 alpha (HIF1A) (NM_001530) Human Tagged Lenti ORF Clone            |
| Tag:                      | Myc-DDK   |
| Symbol:                   | HIF-1 alpha   |
| Synonyms:                 | bHLHe78; HIF-1-alpha; HIF-1A; HIF-1alpha; HIF1; HIF1-ALPHA; MOP1; PASD8 |
| Mammalian Cell Selection: | Puromycin   |
| Vector:                   | pLenti-C-Myc-DDK-P2A-Puro (PS100092)                                    |
| E. coli Selection:        | Chloramphenicol (34 ug/mL)  |
| ORF Nucleotide Sequence:  | The ORF insert of this clone is exactly the same as(RC202461).          |
| Restriction Sites:        | SgfI-MluI   |
| Cloning Scheme:           |   |

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF.


|           |           |
|-----------|-----------|
| ACCN:     | NM_001530 |
| ORF Size: | 2478 bp   |



[View online >](#)

|                               |  |
|-------------------------------|--|
| <b>OTI Disclaimer:</b>        | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>   |
| <b>OTI Annotation:</b>        | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.   |
| <b>Components:</b>            | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).   |
| <b>Reconstitution Method:</b> | <ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>  |
| <b>RefSeq:</b>                | <a href="#">NM_001530.2</a>  |
| <b>RefSeq Size:</b>           | 3958 bp  |
| <b>RefSeq ORF:</b>            | 2481 bp  |
| <b>Locus ID:</b>              | 3091   |
| <b>UniProt ID:</b>            | <a href="#">Q16665</a>   |
| <b>Cytogenetics:</b>          | 14q23.2  |
| <b>Domains:</b>               | PAS, HLH, PAC  |
| <b>Protein Families:</b>      | Transcription Factors  |
| <b>Protein Pathways:</b>      | mTOR signaling pathway, Pathways in cancer, Renal cell carcinoma   |
| <b>MW:</b>                    | 92.5 kDa   |
| <b>Gene Summary:</b>          | This gene encodes the alpha subunit of transcription factor hypoxia-inducible factor-1 (HIF-1), which is a heterodimer composed of an alpha and a beta subunit. HIF-1 functions as a master regulator of cellular and systemic homeostatic response to hypoxia by activating transcription of many genes, including those involved in energy metabolism, angiogenesis, apoptosis, and other genes whose protein products increase oxygen delivery or facilitate metabolic adaptation to hypoxia. HIF-1 thus plays an essential role in embryonic vascularization, tumor angiogenesis and pathophysiology of ischemic disease. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene. [provided by RefSeq, Jul 2011] |

## Product images:



Double digestion of RC202461L3 using SgfI and MluI