

Product datasheet for RC202472L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PKC zeta (PRKCZ) (NM_002744) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: PKC zeta (PRKCZ) (NM_002744) Human Tagged ORF Clone Lentiviral Particle

Symbol: PKC zeta

Synonyms: PKC-ZETA; PKC2

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag:Myc-DDKACCN:NM_002744

ORF Size: 1776 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC202472).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 002744.3

 RefSeq Size:
 2359 bp

 RefSeq ORF:
 1779 bp

 Locus ID:
 5590

 UniProt ID:
 Q05513

 Cytogenetics:
 1p36.33

Domains: PB1, pkinase, S_TK_X, TyrKc, DAG_PE-bind, S_TKc

Protein Families: Druggable Genome, Protein Kinase





PKC zeta (PRKCZ) (NM_002744) Human Tagged ORF Clone Lentiviral Particle - RC202472L3V

Protein Pathways: Chemokine signaling pathway, Endocytosis, Insulin signaling pathway, Tight junction, Type II

diabetes mellitus

MW: 67.7 kDa

Gene Summary: Protein kinase C (PKC) zeta is a member of the PKC family of serine/threonine kinases which

are involved in a variety of cellular processes such as proliferation, differentiation and secretion. Unlike the classical PKC isoenzymes which are calcium-dependent, PKC zeta exhibits a kinase activity which is independent of calcium and diacylglycerol but not of phosphatidylserine. Furthermore, it is insensitive to typical PKC inhibitors and cannot be activated by phorbol ester. Unlike the classical PKC isoenzymes, it has only a single zinc finger module. These structural and biochemical properties indicate that the zeta subspecies is related to, but distinct from other isoenzymes of PKC. Alternative splicing results in multiple

transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008]