

## Product datasheet for RC202476L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Dopamine D2 Receptor (DRD2) (NM\_000795) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type: Lentiviral Particles

Product Name: Dopamine D2 Receptor (DRD2) (NM 000795) Human Tagged ORF Clone Lentiviral Particle

Symbol: DRD2

Synonyms: D2DR; D2R

**Mammalian Cell** 

None

Selection:

**Vector:** pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 000795

ORF Size: 1329 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC202476).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 000795.2</u>

 RefSeq Size:
 2713 bp

 RefSeq ORF:
 1332 bp

 Locus ID:
 1813

 UniProt ID:
 P14416

 Cytogenetics:
 11q23.2

**Protein Families:** Druggable Genome, GPCR, Transmembrane

**Protein Pathways:** Gap junction, Neuroactive ligand-receptor interaction





Dopamine D2 Receptor (DRD2) (NM\_000795) Human Tagged ORF Clone Lentiviral Particle – RC202476L1V

MW: 50.6 kDa

**Gene Summary:** This gene encodes the D2 subtype of the dopamine receptor. This G-protein coupled receptor

inhibits adenylyl cyclase activity. A missense mutation in this gene causes myoclonus dystonia; other mutations have been associated with schizophrenia. Alternative splicing of this gene results in two transcript variants encoding different isoforms. A third variant has been described, but it has not been determined whether this form is normal or due to

aberrant splicing. [provided by RefSeq, Jul 2008]