

## Product datasheet for RC202488L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## BLNK (NM\_013314) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** BLNK (NM\_013314) Human Tagged ORF Clone Lentiviral Particle

Symbol: BLNK

Synonyms: AGM4; BASH; bca; BLNK-S; LY57; SLP-65; SLP65

**Mammalian Cell** 

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK

ACCN: NM\_013314

ORF Size: 1368 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC202488).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 013314.2

 RefSeq Size:
 1829 bp

 RefSeq ORF:
 1371 bp

 Locus ID:
 29760

 UniProt ID:
 Q8WV28

 Cytogenetics:
 10q24.1

**Protein Families:** Druggable Genome

**Protein Pathways:** B cell receptor signaling pathway, Primary immunodeficiency





ORIGENE

**MW:** 50.5 kDa

Gene Summary: This ger

This gene encodes a cytoplasmic linker or adaptor protein that plays a critical role in B cell development. This protein bridges B cell receptor-associated kinase activation with downstream signaling pathways, thereby affecting various biological functions. The phosphorylation of five tyrosine residues is necessary for this protein to nucleate distinct signaling effectors following B cell receptor activation. Mutations in this gene cause hypoglobulinemia and absent B cells, a disease in which the pro- to pre-B-cell transition is developmentally blocked. Deficiency in this protein has also been shown in some cases of pre-B acute lymphoblastic leukemia. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, May 2012]