

## Product datasheet for RC202514L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CHST12 (NM\_018641) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** CHST12 (NM\_018641) Human Tagged ORF Clone Lentiviral Particle

Symbol: CHST12

Synonyms: C4S-2; C4ST-2; C4ST2

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_018641 **ORF Size:** 1242 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC202514).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 018641.3

RefSeq Size: 2159 bp
RefSeq ORF: 1245 bp
Locus ID: 55501
UniProt ID: Q9NRB3
Cytogenetics: 7p22.3

**Domains:** Sulfotransfer2

**Protein Families:** Transmembrane





## CHST12 (NM\_018641) Human Tagged ORF Clone Lentiviral Particle - RC202514L4V

**Protein Pathways:** Chondroitin sulfate biosynthesis, Sulfur metabolism

MW: 48.4 kDa

**Gene Summary:** The protein encoded by this gene belongs to the sulfotransferase 2 family. It is localized to

the golgi membrane, and catalyzes the transfer of sulfate to position 4 of the N-acetylgalactosamine (GalNAc) residue of chondroitin and desulfated dermatan sulfate. Chondroitin sulfate constitutes the predominant proteoglycan present in cartilage, and is distributed on the surfaces of many cells and extracellular matrices. Alternatively spliced transcript variants differing only in their 5' UTRs have been found for this gene. [provided by

RefSeq, Aug 2011]