

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC202529L3V

NDUFAF7 (NM_144736) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	NDUFAF7 (NM_144736) Human Tagged ORF Clone Lentiviral Particle
Symbol:	NDUFAF7
Synonyms:	C2orf56; MidA; PRO1853
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_144736
ORF Size:	1323 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC202529).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 144736.3</u>
RefSeq Size:	2221 bp
RefSeq ORF:	1326 bp
Locus ID:	55471
UniProt ID:	<u>Q7L592</u>
Cytogenetics:	2p22.2
Domains:	DUF185
MW:	49.2 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:This gene encodes an assembly factor protein which helps in the assembly and stabilization
of Complex I, a large multi-subunit enzyme in the mitochondrial respiratory chain. Complex I
is involved in several physiological activities in the cell, including metabolite transport and
ATP synthesis. The encoded protein is a methyltransferase which methylates Arg85 of a
subunit of Complex I in the early stages of its assembly. A pseudogene related to this gene is
located on chromosome 8. Alternative splicing results in multiple transcript variants.
[provided by RefSeq, Sep 2016]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US