


Product datasheet for RC202530L2

RIP2 (RIPK2) (NM_003821) Human Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	RIP2 (RIPK2) (NM_003821) Human Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	RIP2
Synonyms:	CARD3; CARDIAK; CCK; GIG30; RICK; RIP2
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC202530).
Restriction Sites:	Sgfl-Mlul
Cloning Scheme:	
	Cloning sites used for ORF Shuttling:
	Sgf I ORF Mlu I GCG ATC GC ATG // NNN ACG CGT



ACCN: ORF Size: NM_003821 1620 bp

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

varies depending on the nature of the gene.Components:The ORF clone is ion-exchange column purified and shippe containing 10ug of transfection-ready, dried plasmid DNA (Reconstitution Method:1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to 3. Close the tube and incubate for 10 minutes at room tem 4. Briefly vortex the tube and then do a quick spin (less tha at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable shipping when stored at -20°C.RefSeq:NM 003821.4RefSeq ORF:1623 bpLocus ID:043353Oyagabel Cytogenetics:8q21.3Domains:pkinase, TyrKc, CARD, S_TKcProtein Families:Druggable Genome, Protein KinaseProtein Pathways:Neurotrophin signaling pathway, NOD-like receptor signaling MW:	C202530L2	ORÎGENE RIP2 (RII
reference only. However, individual transcript sequences o naturally occurring variations (e.g. polymorphisms), each w clone is substantially in agreement with the reference, but variants is recommended prior to use. More infoOTI Annotation:This clone was engineered to express the complete ORF wi varies depending on the nature of the gene.Components:The ORF clone is ion-exchange column purified and shippe containing 10ug of transfection-ready, dried plasmid DNA (Reconstitution Method:1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to 3. Close the tube and incubate for 10 minutes at room tem 4. Briefly vortex the tube and then do a quick spin (less tha at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable shipping when stored at -20°C.RefSeq:NM 003821.4RefSeq ORF:1623 bpLocus ID:043353Oytogenetics:8q21.3Domains:pkinase, TyrKc, CARD, S_TKcProtein Families:Druggable Genome, Protein KinaseProtein Pathways:Neurotrophin signaling pathway, NOD-like receptor signaliMW:61.2 kDaGene Summary:This gene encodes a member of the receptor-interacting p serine/threonine protein kinases. The encoded protein com activation and recruitment domain (CARD), and is a compo both the innate and adaptive immune pathways. It is a pot	nd/or rearrangements. Therefore, his plasmid DNA. Additional amounts :ific, full-sequence verification at a)TI Disclaimer:
varies depending on the nature of the gene.Components:The ORF clone is ion-exchange column purified and shippe containing 10ug of transfection-ready, dried plasmid DNA (Reconstitution Method:1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to 3. Close the tube and incubate for 10 minutes at room tem 4. Briefly vortex the tube and then do a quick spin (less tha at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable 	s of the same gene can differ through n with its own valid existence. This	
containing 10ug of transfection-ready, dried plasmid DNA (Reconstitution Method:1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to 3. Close the tube and incubate for 10 minutes at room tem 4. Briefly vortex the tube and then do a quick spin (less tha 	with an expression tag. Expression)TI Annotation:
2. Carefully open the tube and add 100ul of sterile water to 3. Close the tube and incubate for 10 minutes at room tem 4. Briefly vortex the tube and then do a quick spin (less tha at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable shipping when stored at -20°C.RefSeq:NM 003821.4RefSeq Size:2588 bpRefSeq ORF:1623 bpLocus ID:8767UniProt ID:O43353Cytogenetics:8q21.3Domains:pkinase, TyrKc, CARD, S_TKcProtein Families:Druggable Genome, Protein KinaseProtein Pathways:Neurotrophin signaling pathway, NOD-like receptor signaliMW:61.2 kDaGene Summary:This gene encodes a member of the receptor-interacting protein kinases. The encoded protein cor activation and recruitment domain (CARD), and is a compo both the innate and adaptive immune pathways. It is a pote	-	omponents:
RefSeq Size:2588 bpRefSeq ORF:1623 bpLocus ID:8767UniProt ID:O43353Cytogenetics:8q21.3Domains:pkinase, TyrKc, CARD, S_TKcProtein Families:Druggable Genome, Protein KinaseProtein Pathways:Neurotrophin signaling pathway, NOD-like receptor signaliMW:61.2 kDaGene Summary:This gene encodes a member of the receptor-interacting protein kinases. The encoded protein corr activation and recruitment domain (CARD), and is a comport both the innate and adaptive immune pathways. It is a potentic corr	emperature. han 5000xg) to concentrate the liquid	econstitution Method:
RefSeq ORF:1623 bpLocus ID:8767UniProt ID:O43353Cytogenetics:8q21.3Domains:pkinase, TyrKc, CARD, S_TKcProtein Families:Druggable Genome, Protein KinaseProtein Pathways:Neurotrophin signaling pathway, NOD-like receptor signaliMW:61.2 kDaGene Summary:This gene encodes a member of the receptor-interacting pathway. The serine/threonine protein kinases. The encoded protein contactivation and recruitment domain (CARD), and is a compo both the innate and adaptive immune pathways. It is a potential in the innate and adaptive immune pathways. It is a potential in the innate and adaptive immune pathways. It is a potential in the innate and adaptive immune pathways. It is a potential in the innate and adaptive immune pathways. It is a potential in the innate and adaptive immune pathways.		efSeq:
Locus ID:8767UniProt ID:O43353Cytogenetics:8q21.3Domains:pkinase, TyrKc, CARD, S_TKcProtein Families:Druggable Genome, Protein KinaseProtein Pathways:Neurotrophin signaling pathway, NOD-like receptor signaliMW:61.2 kDaGene Summary:This gene encodes a member of the receptor-interacting pathway in the innate and adaptive immune pathways. It is a potential in the innate and adaptive immune pathways. It is a potential in the innate and adaptive immune pathways. It is a potential in the innate and adaptive immune pathways. It is a potential in the innate and adaptive immune pathways. It is a potential in the innate and adaptive immune pathways.		efSeq Size:
UniProt ID:O43353Cytogenetics:8q21.3Domains:pkinase, TyrKc, CARD, S_TKcProtein Families:Druggable Genome, Protein KinaseProtein Pathways:Neurotrophin signaling pathway, NOD-like receptor signaliMW:61.2 kDaGene Summary:This gene encodes a member of the receptor-interacting protein kinases. The encoded protein con activation and recruitment domain (CARD), and is a compo both the innate and adaptive immune pathways. It is a potential in the innate and adaptive immune pathways. It is a potential in the innate and adaptive immune pathways.		efSeq ORF:
Cytogenetics:8q21.3Domains:pkinase, TyrKc, CARD, S_TKcProtein Families:Druggable Genome, Protein KinaseProtein Pathways:Neurotrophin signaling pathway, NOD-like receptor signaliMW:61.2 kDaGene Summary:This gene encodes a member of the receptor-interacting pathway is a compo both the innate and adaptive immune pathways. It is a potential is a potential in the innate and adaptive immune pathways. It is a potential is a potential in the innate and adaptive immune pathways.		ocus ID:
Domains:pkinase, TyrKc, CARD, S_TKcProtein Families:Druggable Genome, Protein KinaseProtein Pathways:Neurotrophin signaling pathway, NOD-like receptor signaliMW:61.2 kDaGene Summary:This gene encodes a member of the receptor-interacting protein kinases. The encoded protein con activation and recruitment domain (CARD), and is a compo both the innate and adaptive immune pathways. It is a potential of the second s		IniProt ID:
Protein Families:Druggable Genome, Protein KinaseProtein Pathways:Neurotrophin signaling pathway, NOD-like receptor signaliMW:61.2 kDaGene Summary:This gene encodes a member of the receptor-interacting particular serine/threonine protein kinases. The encoded protein con activation and recruitment domain (CARD), and is a compo both the innate and adaptive immune pathways. It is a potential series of the series o		ytogenetics:
Protein Pathways:Neurotrophin signaling pathway, NOD-like receptor signaliMW:61.2 kDaGene Summary:This gene encodes a member of the receptor-interacting particular serine/threonine protein kinases. The encoded protein con activation and recruitment domain (CARD), and is a compo both the innate and adaptive immune pathways. It is a potential series of the		omains:
MW: 61.2 kDa Gene Summary: This gene encodes a member of the receptor-interacting plaserine/threonine protein kinases. The encoded protein con activation and recruitment domain (CARD), and is a compo both the innate and adaptive immune pathways. It is a potential of the innate and adaptive immune pathways. It is a potential of the innate and adaptive immune pathways.		rotein Families:
Gene Summary: This gene encodes a member of the receptor-interacting portion of the receptor-interacting portion and recruitment kinases. The encoded protein contractivation and recruitment domain (CARD), and is a comport both the innate and adaptive immune pathways. It is a potential of the innate and adaptive immune pathways.	aling pathway	rotein Pathways:
serine/threonine protein kinases. The encoded protein con activation and recruitment domain (CARD), and is a compo both the innate and adaptive immune pathways. It is a pot		/W:
	contains a C-terminal caspase ponent of signaling complexes in potent activator of NF-kappaB and	iene Summary:

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Product images:



Circular map for RC202530L2

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US